

Pharmacy and Drug Development

Research Article

Growth Promoting Potential of Caesalpinia Bonduc Leaf Oil in Broiler Diets: A Natural Alternative to Synthetic Antibiotics

Alagbe Olujimi John^{1*}

¹Department of Animal Production, Gandhi College of Agriculture, Rajasthan India

¹Department of Animal Science, Centre for Distance Learning and Continuous Education, University of Abuja Nigeria

***Corresponding Author:** Alagbe Olujimi John, Department of Animal Production, Gandhi College of Agriculture, Rajasthan India.

Received Date: 16 March 2026; **Accepted Date:** 10 April 2026; **Published Date:** 15 April 2026

Copyright: © 2026 Alagbe Olujimi John, this is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

The global push for antibiotics free poultry production has intensified the search for natural growth promoters. This study evaluated the effects of graded levels of Caesalpinia bonduc leaf oil on the growth performance and nutrient digestibility of broiler chickens. A total of 300 -1 day old broiler chicks were randomly assigned into five dietary treatments in a Completely Randomized Design for a 42 day trial. The birds were reared in a battery cage and fed basal diet formulated to NRC (1994) standards. The treatment consisted of: Treatment 1 (T1) basal diet only; T2 (basal diet with 0.5 mL Caesalpinia bonduc leaf oil/kg diet; T3 (basal diet with 1.0 mL Caesalpinia bonduc leaf oil/kg diet; T4 (basal diet with 1.5 mL Caesalpinia bonduc leaf oil/kg diet and T5 (basal diet with 2.0 mL Caesalpinia bonduc leaf oil/kg diet. Phytochemical analysis of T2 (basal diet with 0.5 mL Caesalpinia bonduc leaf showed that it contained phenols (13.76 %), flavonoids (10.22 %), terpenoids (8.77 %), tannins (4.31 %), and saponins (1.87 %). Result showed that the overall body weight gain was significantly higher ($p < 0.05$) in birds receiving Caesalpinia bonduc leaf oil (T2-T5) compared to the control group (T1). Correspondingly, the feed to gain ratio significantly improved ($p < 0.05$) in oil supplemented groups. Nutrient digestibility of dry matter, organic matter, crude protein, ether extract and ash was significantly enhanced ($p < 0.05$) in birds fed Caesalpinia bonduc leaf oil diet. It was concluded that Caesalpinia bonduc leaf oil at an inclusion of 2.0 mL/kg is an effective phytogetic feed additive for optimizing growth and nutrient utilization in broiler chickens.

Keywords: Caesalpinia bonduc, PhytoGENICS, broiler, Performance, Phytochemicals, Nutrient digestibility.

Introduction

The global poultry industry is currently undergoing a significant transition toward sustainable, antibiotic-free production systems [1,2]. Traditionally, antibiotic growth promoters were used extensively to enhance performance and control subclinical infections; however, their use has been largely restricted or banned in many countries due to the rising threat of antimicrobial resistance and the presence of drug residues in animal products [3,4]. This shift has catalyzed research into phytogetic feed additives, basically derived from plants, such as herbs, spices, and essential oils—as effective natural alternatives [5,6].

Among these natural alternatives, Caesalpinia bonduc (L.) Roxb., commonly known as the "grey nicker nut," stands

out due to its rich profile of secondary metabolites, including tannins, flavonoids, alkaloids, and terpenoids [7,8]. Essential oils extracted from its leaves are particularly valued for their lipophilic nature and broad-spectrum antimicrobial activity [9,10]. These oils have the potential to modulate the intestinal microbiota, stimulate the secretion of endogenous digestive enzymes, and improve the morphology of the gastrointestinal tract, thereby enhancing both growth performance and nutrient utilization in broiler chickens [11,12].

Despite the well-documented potential of various phytoGENICS, the poultry industry faces persistent challenges in maintaining high productivity and bird health in the absence of conventional antibiotics [13,14]. Standard diets often fail to

provide the necessary gut health support needed to maximize feed efficiency and nutrient digestibility, leading to increased susceptibility to enteric pathogens and economic losses [15,16,17].

Furthermore, while *Caesalpinia bonduc* is known for its pharmacological properties, there is a significant lack of standardized data regarding its application as a leaf essential oil in broiler nutrition. The efficacy of phytogenic feed additives can be highly inconsistent due to variations in inclusion levels, plant part used, and the specific chemical composition of the oil [18,19]. Without a comprehensive evaluation of its specific effects on growth parameters and nutrient digestibility, its potential as a reliable feed additive remains largely untapped and unoptimized for commercial use.

This study is justified by the urgent need for safe, residue-free, and cost-effective growth promoters that align with modern consumer demands for safe products from birds [20,21]. *Caesalpinia bonduc* is an abundant, locally available medicinal plant that offers a sustainable alternative to synthetic chemicals [22,23].

Materials And Methods

Experimental site and duration

The experiment was conducted over a 42 days period at the Poultry Section, Gandhi College of Agricultural Technology, Rajasthan, India between the months of June – September 2025. All experimental procedures followed the guidelines for the care and use of experimental birds outlined at the Animal Production and Health Department of the Institution (ASD/008L/2025).

Preparation of plant materials, Identification and extraction of oil

Fresh *Caesalpinia bonduc* leaves were harvested in the early morning within the Biological Sciences Department garden, Gandhi College of Agriculture, Rajasthan India and taken to Botany Department of the same institution for proper authentication by Mr. Ham Singh and assigned a reference number GD/09/2411 for future botanical verification. The collected leaves were manually sorted, washed with distilled and spread on a plastic tray to dry under shade for 12 days until a constant weight was achieved. The dried leaves was pulverized into coarse powder with mortar and pestle to increase the surface area for extraction and stored in an airtight containers until further use. 100 g of pulverized *Caesalpinia bonduc* leaf was taken to the laboratory to determine the composition of phytochemicals in the sample. Analysis (Table 2) was carried out according to the methods outlined by Harbone (1998).

Extraction of *Caesalpinia bonduc* leaf oil was carried out using Clavenger –type equipment. 200 g of pulverized *Caesalpinia bonduc* leaves was placed in a 2 liter round bottom flask mixed with distilled water at a ratio of 1:3 (w/v). The flask is mounted on a heating mantle and maintained at a

temperature of 80 °C for 45 minutes, mixture of water and oil vapor passes through the condenser and it is collected in a graduated receiver. The oil layer which floats on the surface was carefully removed from the mixture residual moisture in the oil was removed by adding 0.5 g of anhydrous sodium sulfate. Final oil was transferred into glass vial and stored in the refrigerator at a temperature of 4°C.

Experimental birds and management practices

300 – 1 day old broiler chicks (Ross 307) was used for this experiment. Upon arrival, birds were unboxed, weighed with automated digital weighing scale to obtain their initial body weight gain and randomly distributed into one of the five dietary treatment groups, each treatment group consisted of 60 birds. The birds were housed in a clean disinfected battery cage measuring 0.8 m by 0.5 m by 0.35 m (L×B×H). Brooding was done in the first two weeks using gas heaters to maintain optimum temperature. Clean drinking water and basal diets were provided throughout the experimental period. Vaccinations (Table 2) were strictly followed according to the Veterinary Regulations in Rajasthan.

Experimental diets and design

A completely randomized design was adopted and basal diet was formulated to meet the specified nutrient requirements of broiler chickens according to NRC (1994) recommendations. The birds were distributed into 5 treatments with 6 replicates consisting of 10 birds each. The treatment group were as follows:

Treatment 1 (control): Basal diet only without *Caesalpinia bonduc* leaf oil

Treatment 2: Basal diet with 0.5 mL + *Caesalpinia bonduc* leaf oil /kg diet

Treatment 3: Basal diet with 1.0 mL + *Caesalpinia bonduc* leaf oil /kg diet

Treatment 4: Basal diet with 1.5 mL + *Caesalpinia bonduc* leaf oil /kg diet

Treatment 5: Basal diet with 2.0 mL + *Caesalpinia bonduc* leaf oil /kg diet

To ensure the full dose of *Caesalpinia bonduc* leaf oil was consumed, daily oil requirements of each group was mixed with 100 g of feed and offered as the first meal in the morning. The remainder of the daily feed allowance was provided after the treated portion was fully consumed.

Data collection

Birds were weighed weekly in the morning before feeding to evaluate the body weight gain. Feed consumed was recorded daily by subtracting the weight of the left over from the weight of feed given. Feed to gain ratio was calculated as the ratio of feed consumed to weight gain.

Nutrient digestibility trial

Nutrient digestibility trial was carried out on the last day of the trial. 5 birds were randomly selected per treatment and

transferred into a metabolic cage. Birds were allowed to adjust to their new environment for 2 days. The digestibility trial lasted for 5 days, a known quantity of feed was given to birds, total excreta were collected, weighed and oven dried at a temperature of 70 °C for 4 hours. Samples of feed and excreta were analyzed for dry matter, crude protein, crude fibre, ether extract and ash using CSD® near infra-red feed analyzer (Model 3400C, China), calibrated and adjusted to the manufacturer's instruction to ensure precision in results

% Nutrient digestibility

$$= \frac{\text{Nutrient intake} - \text{Nutrient voided}}{\text{Nutrient intake}} \times 100$$

Statistical analysis

All data collected on growth performance and nutrient digestibility were subjected to One-way Analysis of Variance for Completely Randomized Design. Significant difference between treatment means were separated using Duncan Multiple Range Test at a probability level of p<0.05.

Ingredients	Diet 1 (Starter diet; 0-21d)	Diet 2 (Finisher diet; 22-42d)
Maize	50.00	53.55
Wheat bran	2.15	2.00
Soya meal	37.00	32.00
Fish meal	2.00	2.00
Limestone	2.50	3.00
Bone meal	5.00	6.00
Methionine	0.25	0.25
Lysine	0.25	0.25
*Mineral-Vitamin Premix	0.25	0.25
Sodium chloride	0.40	0.50
Toxin binderTMR	0.20	0.20
Total	100.00	100.00
Determined analysis (%)		
Dry matter	87.12	89.36
Crude protein	23.16	21.09
Crude fibre	3.98	4.11
Ether extract	3.53	3.95

Calcium	1.73	1.85
Phosphorus	0.82	0.87
Energy (Kcal/kg)	3009.5	3201.7

Table 1: Ingredient and chemical composition of basal diet (% DM)

Starter Min-Vitamin Premix; 2.5 kg contains—vitamin A: 10,000,000 IU; vitamin D3: 4,000,000 IU; vitamin E: 27,000 IU; vitamin K: 3,000 mg; thiamine B1: 1,800 mg; riboflavin B2: 7000 mg; pyridoxine B6: 2,800 mg; niacin: 30,500 mg; vitamin B12: 20 mg; pantothenic acid: 9000 mg; folic acid: 8000 mg; biotin: 70 mg; choline chloride: 350 g; antioxidant: 115 mg; magnesium: 100 g; zinc: 70 g; iron: 13 g; copper: 8.00 g; iodine: 1.25 g; selenium: 100 mg; cobalt: 100 mg. Finisher Min-Vitamin Premix; 2.5 kg contains —vitamin A: 10,000,000 IU; vitamin D3: 2,000,000 IU; vitamin E: 20,000 IU; vitamin K: 2,250 mg; thiamine B1: 1750 mg; riboflavin B2: 5000 mg; pyridoxine B6: 2750 mg; niacin: 27,500 mg; vitamin B12: 15 mg; pantothenic acid: 7500 mg; folic acid: 7500 mg; biotin: 50 mg; choline chloride: 400 g; antioxidant: 125 mg; magnesium: 80 g; zinc: 50 g; iron: 20 g; copper: 5 g; iodine: 1.2 g; selenium: 200 mg; cobalt: 200 mg.

Days	Vaccines	Route of administration
1	Marek's (at hatchery)	Subcutaneous
7	RDV F1	Ocular
11	RDV Lasota	Drinking water
16	IBD	Drinking water
21	Lasota (booster)	Drinking water
27	IBD (booster)	Drinking water

Table 2: Vaccination schedule for birds

Components	% Composition
Tannins	4.31
Flavonoids	10.22
Alkaloids	3.55
Saponins	1.87
Phenols	13.76
Terpenoids	8.77

Table 3: Phytochemical composition Caesalpinia bonduc leaf

Starter phase (0 - 21d)							
Parameters	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	SEM	p-value
Initial body weight (g/bird)	47.87	48.01	47.66	47.89	47.93	0.02	0.07
Final body weight (g/bird)	859.9 ^b	1061.2 ^a	1066.4 ^a	1068.4 ^a	1069.6 ^a	45.12	0.02
Body weight gain (g/bird)	812.03 ^b	1013.19 ^a	1018.74 ^a	1020.51 ^a	1021.67 ^a	40.80	0.01

Daily weight gain (g/bird)	38.66 ^b	48.25 ^a	48.51 ^a	48.59 ^a	48.65 ^a	0.96	0.001
Feed consumption (g/bird)	1748.2 ^b	1954.2 ^a	1960.9 ^a	1961.2 ^a	1963.2 ^a	61.98	0.03
Daily consumption (g/bird)	83.24 ^b	93.05 ^a	93.37 ^a	93.39 ^a	93.48 ^a	0.18	0.01
Feed to gain ratio	2.15 ^a	2.00 ^b	2.00 ^b	2.00 ^b	2.00 ^b	0.01	0.001
Finisher phase (22-42 d)							
Body weight gain (g/bird)	1441.6 ^b	1702.1 ^a	1730.2 ^a	1735.1 ^a	1741.4 ^a	44.86	0.02
Daily weight gain (g/bird)	68.64 ^b	81.05 ^a	82.39 ^a	82.62 ^a	82.92 ^a	2.65	0.001
Feed consumption (g/bird)	3100.0 ^b	3239.0 ^a	3249.1 ^a	3251.8 ^a	3258.1 ^a	136.8	0.04
Daily feed consumption (g/bird)	147.6 ^b	154.2 ^a	154.7 ^a	154.8 ^a	155.1 ^a	0.44	0.21
Feed to gain ratio	2.15 ^a	2.00 ^b	2.00 ^b	2.00 ^b	2.00 ^b	0.01	0.001
Overall production (1- 42d)							
Body weight gain (g/bird)	2253.63 ^b	2715.29 ^a	2748.94 ^a	2755.61 ^a	2763.07 ^a	103.6	0.03
Daily weight gain (g/bird)	53.65 ^b	64.64 ^a	65.45 ^a	65.60 ^a	65.78 ^a	2.56	0.001
Total feed consumption (g/bird)	4848.2 ^b	5193.2 ^a	5210.0 ^a	5213.0 ^a	5221.3 ^a	0.81	0.001
Daily feed consumption (g/bird)	115.4 ^b	123.6 ^a	124.0 ^a	124.1 ^a	124.3 ^a	0.17	0.006
Feed to gain ratio	2.17 ^a	2.00 ^b	2.00 ^b	2.00 ^b	1.95 ^a	0.11	0.01

^{a,b,c} Means within the same row with different superscript differ significantly ($p < 0.05$)

Table 4: Growth performance of broiler chickens fed different levels of *Caesalpinia bonduc* leaf oil

Components (%)	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5		
Dry matter	79.17 ^b	83.12 ^a	83.88 ^a	84.02 ^a	84.05 ^a	SEM	P-value
Organic matter	38.74 ^b	45.28 ^a	46.77 ^a	46.93 ^a	46.98 ^a	2.63	0.01
Crude protein	69.06 ^b	76.33 ^a	77.82 ^a	77.93 ^a	77.95 ^a	4.08	0.001
Crude fibre	57.08 ^a	48.11 ^b	48.83 ^b	48.86 ^b	48.91 ^b	1.90	0.004
Ether extract	61.94 ^b	72.01 ^a	73.22 ^a	73.25 ^a	73.28 ^a	0.56	0.0001
Ash	55.35 ^b	61.94 ^a	62.88 ^a	63.04 ^a	63.21 ^a	0.42	0.02

^{a,b,c} Means within the same row with different superscript differ significantly ($p < 0.05$)

Table 5: Nutrient digestibility of broiler chickens fed different levels of *Caesalpinia bonduc* leaf oil

Results and Discussion

Table 4 reveals the growth performance of broiler chickens fed different levels of *Caesalpinia bonduc* leaf oil at starter and finisher phase. Variables obtained in this study follow similar trend as body weight gain, feed to gain ratio and feed consumption were significantly ($p < 0.05$) influenced by the treatment. In the overall performance, body weight gain was higher in T2-T5 compared to T1. Total feed consumed was lower in T1 (4848.2 g/bird) than T2 (5193.2 g/b), T3 (5210.0 g/b), T4 (5213.0 g/b) and T5 (5221.3 g/b). Feed to gain ratio was higher in T1 compared to the other treatments.

The observed increase in average daily body weight and significant reduction in feed conversion ratio in treatments T2 – T5 compared to the control (T1) from the starter, finish-

er and overall production indicates that *Caesalpinia bonduc* leaf oil acts as a natural growth promoter. This improvement is likely driven by the diverse phytochemical profile of the essential oil. The high concentration of phenolic compounds (13.76 %) and flavonoids (10.22 %) facilitates the protection of the intestinal mucosal of birds from oxidative stress and the improvement in their overall health [24,25]. Terpenoids (8.77 %) and alkaloids (3.55 %) possess antimicrobial and antibacterial properties that can inhibit pathogenic gut microflora thereby making more energy available for the growth of birds [6,26]. At low concentration, saponins can increase the permeability of intestinal mucosal cells, facilitating efficient nutrient absorption [27]. The final body weight gain obtained

in this study was within 1923.3 – 3000.2 g reported by [28], when phytochemicals was supplemented in the diet of broiler chickens at 3 mL/liter of water. [28] also recorded a feed conversion ratio of 2.1 – 2.4 when herbs at 3 g/kg was added in the diet of broilers. Higher feed consumption all through the starter and finisher phase in birds fed graded levels of *Caesalpinia bonduca* leaf oil indicates that it has an appetizer effect. According to [29], phytochemical compounds that are rich in terpenoids and phenolic compounds often act as sensory stimulators. These compounds provides a unique aroma and flavor to the diet that can stimulate bird's appetite [4]. Although birds have fewer taste buds compared to mammals. However, their olfactory system responds positively to certain plant phyto-constituents leading to increased feed intake [30,31] This outcome suggests that *Caesalpinia bonduca* leaf oil relatively promote balanced intestinal flora thus preventing intestinal inflammation and promoting a healthier gut. Total overall feed consumption range obtained in this study 4848.2 – 5221.3 g/b was similar to 4800.2 g – 5200.8 g reported by [32] when *Polyalthia longifolia* leaf meal was supplemented in the diet of broiler chickens. However, values were lower than 5307.1 – 5600.2 g/b recorded by [33-35], when graded levels of *Eucalyptus* oil was added in the diet of broilers. This variation could be due to concentration of phyto-compounds as well as dose administered. Nutrient digestibility of broiler chickens fed different levels of *Caesalpinia bonduca* leaf oil is revealed in Table 5. Dry matter digestibility value varied from 79.17 - 84.05 %, organic matter (38.74 - 46.98 %), crude protein (69.06 - 77.95 %), crude fibre (48.91 - 57.08 %), ether extract (61.94 - 73.28 %) and Ash (55.35 - 63.21 %). All values were significantly ($p < 0.05$) affected by the treatment. The improved nutrient digestibility of dry matter, organic matter, crude protein and ether extracts in group supplemented with *Caesalpinia bonduca* leaf oil (T2 – T5) explains the physiological basis for lower feed conversion ratio. According to (Shittu et al., 2021), addition of essential oil in the diet of birds can stimulate the activities of endogenous enzymes such as lipase, amylase and trypsin which directly enhances the breakdown and absorption of nutrients [36,37].

Conclusion

The inclusion of *Caesalpinia bonduca* leaf oil at up to 2.0 mL/kg diet successful improved the growth of birds over the 42 days period. The increase in nutritive digestibility suggests that the phyto-components (particularly terpenoids and phenols) enhanced enzymatic activities and intestinal health.

References

1. Abdelli, Nedra, David Solà-Oriol, and José Francisco Pérez. "Phytochemical feed additives in poultry: achievements, prospective and challenges." *Animals* 11, no. 12 (2021): 3471.

2. El-Sayed, A., Fayed, R. H., Castañeda Vázquez, H., & Rügge, K. (2022). Future trend to replace chemical products with nutraceutical food/feed additive: A mini review. *Journal of Applied Veterinary Sciences*, 7(3), 20–29.
3. Alagbe, John Olujimi, Muritala Daniel Shittu, Aduragbemi Yetunde Adesina, Chesa Jummai Grace, Kadiri Mercy Cincinsoke, Bamigboye Samson Oluwafemi, and Effiong Erikano-bong. "The approximate mineral and phytochemical content of the leaves, stem bark, and roots of *Pterocarpus erinaceus* in India." *Cerrado: Agricultural and Biological Research* 1, no. 1 (2024): 32-41.
4. Adewale, A. O., J. O. Alagbe, and Adekemi O. Adeoye. "Dietary supplementation of *Rauvolfia vomitoria* root extract as a phytochemical feed additive in growing rabbit diets: Haematology and serum biochemical indices." *International Journal of Orange Technologies* 3, no. 3 (2021): 1-12.
5. Haque, Md Hakimul, Subir Sarker, Md Shariful Islam, Md Aminul Islam, Md Rezaul Karim, Mohammad Enamul Hoque Kayesh, Muhammad JA Shiddiky, and M. Sawkat Anwer. "Sustainable antibiotic-free broiler meat production: Current trends, challenges, and possibilities in a developing country perspective." *Biology* 9, no. 11 (2020): 411.
6. Ojediran, Taiwo Kayode, I. A. Emiola, V. Durojave, and John Olujimi Alagbe. "Proximate, vitamin and GC-MS profiling of *Kigelia africana* fruit powder." *Cerrado: Agricultural and Biological Research* 1, no. 1 (2024): 13-20.
7. Paraskeuas, Vasileios, Konstantinos Fegeros, Irida Palamidi, Christine Hunger, and Konstantinos C. Mountzouris. "Growth performance, nutrient digestibility, antioxidant capacity, blood biochemical biomarkers and cytokines expression in broiler chickens fed different phytochemical levels." *Animal Nutrition* 3, no. 2 (2017): 114-120.
8. Ri, Chang-Song, Xian-Ren Jiang, Myong-Ho Kim, Jing Wang, Hai-Jun Zhang, Shu-Geng Wu, Valentino Bontempo, and Guang-Hai Qi. "Effects of dietary oregano powder supplementation on the growth performance, antioxidant status and meat quality of broiler chicks." *Italian Journal of Animal Science* 16, no. 2 (2017): 246-252.
9. Sharma, Singh, Alagbe Olujimi John, Liu Xing, Sharma Ram, and Kumar Amita. "Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroemetry." *International Journal of Agriculture and Animal Production* 2, no. 6 (2022): 18-27.
10. Mohiti-Asli, Maziar, and Moein Ghanaatparast-Rashti. "Comparison of the effect of two phytochemical compounds on growth performance and immune response of broilers." *Journal of Applied Animal Research* 45, no. 1 (2017): 603-608.
11. Engberg, Ricarda Margarete, Kai Grevsen, Elise Ivarsen, Xavier Fretté, Lars Porskjær Christensen, Ole Højberg, Bent Borg Jensen, and Nuria Canibe. "The effect of *Artemisia annua* on broiler performance, on intestinal microbiota and on the course of a *Clostridium perfringens* infection applying a necrotic enteritis disease model." *Avian Pathology* 41, no. 4 (2012): 369-376.

12. Giannenas, Ilias, Eleftherios Bonos, Ioannis Skoufos, Athina Tzora, Ioanna Stylianaki, Diamanto Lazari, Anastasios Tsinas, Efterpi Christaki, and Panagiota Florou-Paneri. "Effect of herbal feed additives on performance parameters, intestinal microbiota, intestinal morphology and meat lipid oxidation of broiler chickens." *British Poultry Science* 59, no. 5 (2018): 545-553.
13. Bozkurt, Mehmet, Kamil Küçükyılmaz, Abdullah Uğur Çatlı, Zafer Özyıldız, Mustafa Çınar, Metin Çabuk, and Fethiye Çöven. "Influences of an essential oil mixture supplementation to corn versus wheat-based practical diets on growth, organ size, intestinal morphology and immune response of male and female broilers." *Italian Journal of Animal Science* 11, no. 3 (2012): e54.
14. Bravo, D., P. Utterback, and C. M. Parsons. "Evaluation of a mixture of carvacrol, cinnamaldehyde, and capsicum oleoresin for improving growth performance and metabolizable energy in broiler chicks fed corn and soybean meal." *Journal of Applied Poultry Research* 20, no. 2 (2011): 115-120.
15. Alp, M., Mustafa Midilli, Neşe Kocabağlı, Hüseyin Yılmaz, N. Turan, A. Gargılı, and N. Acar. "The effects of dietary oregano essential oil on live performance, carcass yield, serum immunoglobulin G level, and oocyst count in broilers." *Journal of Applied Poultry Research* 21, no. 3 (2012): 630-636.
16. Amerah, Ahmed M., Alexandre Péron, Fifi Zaefarian, and Velmurugu Ravindran. "Influence of whole wheat inclusion and a blend of essential oils on the performance, nutrient utilisation, digestive tract development and ileal microbiota profile of broiler chickens." *British poultry science* 52, no. 1 (2011): 124-132.
17. Morris, Hernández, and Alagbe John Olujimi. "INFLUENCE OF Odontonema Strictum OIL ON THE GROWTH PERFORMANCE AND RUMINAL FERMENTATION OF BARBARI BUCKS." *Research in: Agricultural & Veterinary Sciences* 9, no. 2 (2025).
18. Isabel, B. A. S. Y., and Y. Santos. "Effects of dietary organic acids and essential oils on growth performance and carcass characteristics of broiler chickens." *Journal of Applied Poultry Research* 18, no. 3 (2009): 472-476.
19. Xu, Z. R., C. H. Hu, M. S. Xia, X. A. Zhan, and M. Q. Wang. "Effects of dietary fructooligosaccharide on digestive enzyme activities, intestinal microflora and morphology of male broilers." *Poultry science* 82, no. 6 (2003): 1030-1036.
20. Hamed, Somayeh, Tahoorah Shomali, and Hossein Ghaderi. "Effect of dietary inclusion of Mentha piperita L. on histological and histomorphometrical parameters of the small intestine in broiler chickens." *Organic Agriculture* 7, no. 2 (2017): 105-110.
21. Park, J. H., and I. H. Kim. "Effects of a protease and essential oils on growth performance, blood cell profiles, nutrient retention, ileal microbiota, excreta gas emission, and breast meat quality in broiler chicks." *Poultry science* 97, no. 8 (2018): 2854-2860.
22. Jamroz, Dariusz, Tomasz Wartecki, Maciej Houszka, and C. Kamel. "Influence of diet type on the inclusion of plant origin active substances on morphological and histochemical characteristics of the stomach and jejunum walls in chicken." *Journal of Animal Physiology and Animal Nutrition* 90, no. 5-6 (2006): 255-268.
23. Chen, Yan, Jun Wang, Longfei Yu, Tianyue Xu, and Nianhua Zhu. "Microbiota and metabolome responses in the cecum and serum of broiler chickens fed with plant essential oils or virginiamycin." *Scientific reports* 10, no. 1 (2020): 5382.
24. Alagbe, J. O. "Dietary supplementation of rauwolfia vomitoria root extract as a phyto-genic feed additive in growing rabbit diets: growth performance and caecal microbial population." *Concept in Dairy and Veterinary Sciences* 4, no. 2 (2021): 2021.
25. İpçak, Hasan Hüseyin. "The role of phyto-genic feed additives in modulating poultry nutritional physiology and genomics." In *Veterinary Medicine and Science Annual Volume 2025*. IntechOpen, 2023.
26. Abdel-Wareth, Ahmed AA, Saskia Kehraus, and Karl-Heinz Südekum. "Peppermint and its respective active component in diets of broiler chickens: growth performance, viability, economics, meat physicochemical properties, and carcass characteristics." *Poultry science* 98, no. 9 (2019): 3850-3859.
27. Noirot, V., R. Moncoulon, D. Sauvant, and C. Bayourthe. "Effect of essential oils and essential oils compounds supplementations in ruminant species: *statistical analysis*." (2007): 589-597.
28. Bashir, Musa, John O. Alagbe, Adegbite Motunrade Betty, and E. A. Omokore. "Growth Performance, caeca microbial population and immune response of starter broiler chicks fed aqueous extract of Balanites aegyptiaca and Alchornea cordifolia stem bark mixture." *United Journal for Research and Technology* 2, no. 2 (2020): 13-21.
29. Alagbe, J. O., M. D. Shittu, and Eunice Abidemi Ojo. "Prospect of leaf extracts on the performance and blood profile of monogastric—A review." *International Journal on Integrated Education* 3, no. 7 (2020): 122-127.
30. Ghazaghi, Mahmoud, Mehran Mehri, and Farzad Bagherzadeh-Kasmani. "Effects of dietary Mentha spicata on performance, blood metabolites, meat quality and microbial ecosystem of small intestine in growing Japanese quail." *Animal Feed Science and Technology* 194 (2014): 89-98.
31. Jang, I. S., Y. H. Ko, S. Y. Kang, and C. Y. Lee. "Effect of a commercial essential oil on growth performance, digestive enzyme activity and intestinal microflora population in broiler chickens." *Animal Feed Science and Technology* 134, no. 3-4 (2007): 304-315.
32. Alagbe, J. O., D. O. Adejumo, S. G. Ademola, A. O. Abiola, B. O. Samson, and F. T. Ushie. "Productive performance, caeca microbial population and immune-modulatory activity of broiler chicks fed different levels Sida acuta leaf extract in replacement of antibiotics." *Bioinformatics and Proteomics Open Access Journal* 5, no. 1 (2021): 000143.

33. Alagbe, John Olujimi. "Effect on performance, serum biochemistry and haematological components of feeding "japanese quails" phytogenic feed additions comprising Megaphrynium macrostachyum leaves." *Brazilian Journal of Science* 3, no. 5 (2024): 51-64.
34. Alagbe J. Olujimi, (2024), Proximate, Mineral Composition and Phyto-Constituents of Some Medicinal Plants/Herbs In India, *J. Pharmaceutics and Pharmacology Research*, 7(12); DOI:10.31579/2688-7517/219
35. Hernandez, M and Alagbe, J.O. (2025a). Influence of Abrus procatorious crude oil supplementation on the growth performance, Nutrient digestibility, Ruminal fermentation and Microbial population of Malabari Bucks. *International Journal of Global Sustainable Research*, 3(7): 527-538.
36. Xu, J., F. Zhou, B-P. Ji, R-S. Pei, and N. Xu. "The antibacterial mechanism of carvacrol and thymol against Escherichia coli." *Letters in applied microbiology* 47, no. 3 (2008): 174-179.
37. Lambert, R. J. W., Proteus N. Skandamis, Proteus J. Coote, and G-JE Nychas. "A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol." *Journal of applied microbiology* 91, no. 3 (2001): 453-462.
38. National Research Council. "Mineral tolerance of animals: 2005." National Research Council. 2005. Mineral Tolerance of Animals: Second Revised Edition (2005): 11309.
39. Haroen, Ucop, Syafwan Syafwan, Kiki Kurniawan, and Agus Budiansyah. "Determination of total phenolics, flavonoids, and testing of antioxidant and antibacterial activities of red ginger (Zingiber officinale var. Rubrum)." *Journal of Advanced Veterinary and Animal Research* 11, no. 1 (2024): 114-124.
40. Attia, Youssef, Mohammed Al-Harhi, and Mahmoud El-Kelawy. "Utilisation of essential oils as a natural growth promoter for broiler chickens." *Italian Journal of Animal Science* 18, no. 1 (2019): 1005-1012.
41. Alagbe, J. O., Adekemi Adeoye, and A. O. Oluwatobi. "Proximate and mineral analysis of Delonix regia leaves and roots." *International Journal on Integrated Education* 3, no. 10 (2020): 144-149.
42. Alagbe, J. O., R. Sharma, Eunice Abidemi Ojo, M. D. Shittu, and Bello Kamoru Atanda. "Chemical evaluation of the proximate, minerals, vitamins and phytochemical analysis of Danielle oliveri stem bark." *International Journal of Biological, Physical and Chemical Studies* 2, no. 1 (2020): 16-22.
43. Alagbe, John Olujimi. "Clerodendron splendens leaf extract supplementation in weaner rabbits: impact on growth performance, haematology and intestinal microbial population." *Cerrado: Agricultural and Biological Research* 1, no. 1 (2024): 21-31.
44. Alagbe, J. O., E. U. Anaso, H. Zubairu, and D. N. Anorue. "Lepadenia Hastate and Cedrus Brevifolia Oil: Effect on Growth Performance, Caecal Microbial Population, Fermentation and Haematological Indices of Weaner Rabbits." *International Journal of Advanced Technology and Social Sciences* 3, no. 9 (2025): 1213-1228.
45. Alagbe Olujimi John (2024). Bioactive profiling of essential oil of Terminalia arjuna stem bark collected from Orathur village, Tamilnadu, india. *Journal of Food Science and Biotechnology*, 1(1): 1-4.