

Skin and Dermatological Research

Research Article

Prevalence and Risk Factors of Acne Vulgaris Among University Students: A Cross-Sectional Study

FATIMA MAZHAR¹, Muhammad Waqar Mazhar^{2*}

¹Department of Microbiology, Muhammad Nawaz Sharif University of Agriculture, Multan, Multan, Pakistan

²Department of Bioinformatics and Biotechnology, Government College University Faisalabad, Pakistan

***Corresponding Author:** Muhammad Waqar Mazhar, Department of Bioinformatics and Biotechnology, Government College University Faisalabad, Pakistan.

Received Date: 15 June 2026; **Accepted Date:** 06 July 2026; **Published Date:** 10 July 2026

Copyright: © 2026 Muhammad Waqar Mazhar, this is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Acne vulgaris is one of the most common skin problems in young people, and it has a big effect on their mental and social lives. The goal is to find out how common acne vulgaris is among college students and what risk factors are linked to it.

Methods: A structured questionnaire and clinical exam were used to do a cross-sectional study with 250 students.

Results: It was found in 68% of people who had acne vulgaris. High amounts of stress ($p < 0.001$), eating a lot of oily foods ($p = 0.002$), and not getting enough sleep ($p = 0.005$) were all important risk factors.

Conclusion: Acne vulgaris is very common among college students and is strongly linked to living factors that can be changed.

Keywords: Acne vulgaris, Dermatology, Prevalence, Risk factors

Introduction

Acne vulgaris is an ongoing, multifactorial inflammatory disorder of the pilosebaceous unit, predominantly impacting teenagers and young adults, however it may continue into later life stages [1]. It is clinically defined by a range of lesions, encompassing open and closed comedones, inflammatory papules, pustules, nodules, and cysts, primarily located on the face, chest, and back. Acne, with a global incidence affecting approximately 80–90% of teenagers, constitutes a major public health issue and significantly contributes to dermatological morbidity globally [2]. The pathophysiology of acne vulgaris is intricate and encompasses four principal processes: heightened sebum production induced by androgens, aberrant follicular keratinisation resulting in comedone formation, colonisation of the pilosebaceous unit by *Cutibacterium acnes*, and ensuing inflammatory responses facilitated by both innate and adaptive immune mechanisms [3]. Recent advancements have underscored the significance of

genetic predisposition, hormonal variations, and modifications in the skin microbiome in affecting disease susceptibility and severity [4]. Moreover, recent evidence indicates that systemic variables, like insulin resistance and high glycaemic load diets, may intensify acne via hormonal and inflammatory mechanisms. Alongside biological variables, lifestyle and environmental factors have garnered heightened focus in recent years [5]. Psychological stress has been demonstrated to aggravate acne through neuroendocrine pathways, specifically via the hypothalamic–pituitary–adrenal (HPA) axis, which can elevate sebum production and inflammatory reactions [6].

Likewise, dietary patterns—particularly elevated consumption of dairy products, fast foods, and high-glycemic-index diets—have been correlated with acne severity in numerous epidemiological studies. Sleep deprivation, sedentary behaviour, and improper cosmetic application exacerbate

acne, especially in university students who frequently endure erratic schedules and academic pressure [7]. Aside from its physical manifestations, acne vulgaris has significant psychosocial consequences. It has been persistently linked to diminished self-esteem, social withdrawal, anxiety, and depression, consequently substantially reducing quality of life. The psychological impact of acne can, in certain instances, be akin to that of chronic systemic diseases, highlighting the necessity for holistic management strategies that encompass both physical and mental health dimensions[8]. Despite the wide global literature on acne, there is a scarcity of region-specific data in developing nations, like Pakistan, where social, nutritional, and environmental factors may significantly differ from those in Western populations. University students constitute a notably susceptible demographic owing to heightened stress exposure, nutritional inconsistencies, and lifestyle alterations [9]. Nevertheless, few research have investigated the synergistic impact of these modifiable risk variables within this demographic. This study is to evaluate the prevalence of acne vulgaris and determine its associated lifestyle and behavioural risk factors among university students. This study aims to enhance the existing knowledge base by offering localised evidence and to aid in the formulation of tailored preventative strategies and educational interventions to alleviate the prevalence of acne among young adults.

Methodology

Study Design and Setting

A cross-sectional analytical investigation was performed during three months (January to March 2026) at a university campus in Pakistan. The study aimed to evaluate the prevalence of acne vulgaris and identify related lifestyle risk factors among university students.

Study Population and Sampling

The study population consisted of undergraduate and postgraduate students aged 18 to 30 years. A sample size of 250 individuals was determined utilising standard prevalence estimation techniques, with a confidence level of 95%, a margin of error of 5%, and an anticipated prevalence of 50% owing to insufficient prior regional data. Participants were selected through a non-probability convenience sampling method from several academic fields.

Inclusion and Exclusion Criteria

Participants aged 18 to 30 years who consented to participate were included in the study. Individuals with chronic dermatological problems excluding acne (e.g., eczema, psoriasis), those undergoing systemic treatment for acne (e.g., isotretinoin, oral antibiotics), and those with recognised endocrine abnormalities impacting skin health were excluded to prevent confounding effects.

Data Collection Procedure

Data were gathered with a standardised, self-administered questionnaire created following an extensive examination of pertinent literature. The questionnaire comprised several sections, encompassing demographic variables (age, gender, academic year), lifestyle factors (dietary habits, sleep length, physical activity), psychological stress levels, and skincare practices (cosmetic usage, frequency of face washing, types of products). Stress levels were evaluated with a validated perceived stress scale or a Likert-type assessment. Before data collection, the questionnaire underwent pilot testing with a small cohort of students (n=20) to ascertain clarity and reliability.

Clinical Assessment

Participants underwent clinical assessment for the existence and severity of acne vulgaris. The diagnosis was established according to standard dermatological criteria, and the severity was assessed utilising the Global Acne Grading System (GAGS), which takes into account lesion kind and anatomical location. According to the scoring system, acne severity was classified as mild, moderate, severe, or extremely severe.

Study Variables

The dependent variable was the occurrence and intensity of acne vulgaris. The independent variables encompassed demographic parameters (age, gender), lifestyle aspects (diet, sleep length, physical activity), psychological stress, and cosmetic usage. Dietary practices primarily centred on the frequency of intake of fatty foods, dairy items, and high-glycemic-index foods.

Statistical Analysis

Data were inputted and analysed utilising Statistical Package for the Social Sciences (SPSS) version 25. Descriptive statistics were computed, encompassing means, standard deviations, frequencies, and percentages. The chi-square test was utilised to evaluate relationships among categorical variables, whereas independent t-tests were employed for continuous variables when applicable. A multivariate logistic regression analysis was conducted to find independent predictors of acne vulgaris, while adjusting for relevant confounders. A p-value below 0.05 was deemed statistically significant.

Ethical Considerations

Before any data was collected, written informed consent was obtained from all subjects. The people who took part were promised privacy and anonymity, and their information was only used for study. All methods were carried out according to the rules and guidelines that were in place.

Results

There were 250 college students who took part in the study,

which means that 100% of them were asked to do so. They were between the ages of 18 and 30 years old, with a mean age of 21.4 ± 2.3 years. 150 of the people who took part were women (60%), and 100 were men (40%).

Variable	Value
Total Participants	250
Mean Age (years)	21.4 ± 2.3
Female (%)	60%
Male (%)	40%

Prevalence of Acne Vulgaris

Out of the whole group, 170 people were clinically diagnosed with acne vulgaris, giving the general prevalence of 68%. The rate was slightly higher in women (72% vs. 62%) than in men, but the difference wasn't statistically significant ($p > 0.05$).

Category	Percentage
Prevalence	68
Mild	40
Moderate	35
Severe	25

Severity of Acne

Using the Global Acne Grading System (GAGS), 170 people with acne were put into three groups based on how bad their acne was: 68 had mild acne (40%), 60 experienced moderate acne (35%), and 42 had serious acne (25%). The people in the study did not have any cases of very bad acne.


Factor	Acne (%)	p-value
High Stress	75	<0.001
Oily Food	70	0.002
Poor Sleep	65	0.005
Cosmetic Use	72	0.01

Association with Lifestyle Factors

A study of lifestyle-related risk factors showed strong links between acne vulgaris and a number of variables that can be changed. 75% of people who said they had high levels of psychological stress had acne, compared to 55% of people who said they had low levels of stress (χ^2 test, $p < 0.001$). In the same way, eating a lot of fat and fast foods was strongly linked to acne, with 70% of people who ate a lot of these foods getting acne compared to only 50% of people who ate less of them ($p = 0.002$).

Predictor	AOR	p-value
Stress	2.4	<0.001

Oily Food	1.9	0.003
Sleep	1.7	0.01
Cosmetics	1.6	0.04



The amount of sleep also had a big effect on the amount of acne. People who said they slept less than 6 hours a night were more likely to have acne (65%) than people who said they slept 6 to 8 hours a night (52%). There was a strong link between cosmetic use and acne, especially among female subjects (72% of cosmetic users had acne compared to 58% of non-users; $p = 0.01$).

Multivariate Analysis

A multivariate logistic regression analysis was done to find independent indicators of acne vulgaris. AOR = 2.4 (95% CI: 1.5–3.8; $p < 0.001$) showed that high psychological stress was a strong independent predictor of acne. Also, eating a lot of fat foods (AOR = 1.9; 95% CI: 1.2–3.0; $p = 0.003$) and not getting enough sleep (AOR = 1.7; 95% CI: 1.1–2.6; $p = 0.01$) were both strong predictors. AOR = 1.6; 95% CI: 1.0–2.5; $p = 0.04$) showed a weak but significant link between cosmetic use and acne.

Additional Findings

There was no statistically significant link ($p > 0.05$) between the number of people with acne and their age or the school year. There was a weak negative relationship between physical exercise and the prevalence of acne, but it wasn't strong enough to be statistically significant ($p = 0.08$).

Discussion

This study looked at how common acne vulgaris is among college students and what living factors may make it worse. It found that 68% of students had acne vulgaris. There are estimates around the world that say between 80% and 90% of teens and young adults have acne. This result supports

those estimates and shows that acne vulgaris is still a very common skin condition in this age group. The slightly higher prevalence seen in female participants compared to males is in line with what other studies have found. These studies show that hormonal changes, cosmetic use, and more health-seeking behaviour in women may make them more vulnerable [10]. One of the most important things this study showed was that there is a strong link between mental worry and acne vulgaris. People who said they were under a lot of stress had a lot more acne, and worry also showed up as an independent predictor in multivariate analysis. This result is supported by research that shows stress can make acne worse by activating the hypothalamic–pituitary–adrenal (HPA) axis and increasing the release of cortisol and androgens. Because these changes in hormones make sebum glands work harder and cause inflammation, acne gets worse [6].

It was also discovered that what you eat has a big impact on getting acne. The study found that eating a lot of oily and fast foods was significantly linked to a higher frequency of acne. These results are in line with earlier study that suggests high-fat and high-glycemic-index diets may raise insulin and insulin-like growth factor-1 (IGF-1) levels, which then cause sebum production and keratinocyte proliferation [11]. The connection between food and acne has been debated in the past, but more and more evidence suggests that changing your diet could be a useful way to treat acne.

The amount of time spent sleeping became another important factor linked to acne vulgaris. Acne was much more common among people who said they didn't get enough sleep (less than six hours per night). Lack of sleep is known to throw off the balance of hormones, raise stress levels, and weaken the immune system. All of these things may make acne worse or cause it to appear in the first place [12]. As part of a complete plan to avoid acne, this result shows how important it is to keep good sleep hygiene.

Cosmetic use, especially among female subjects, was also strongly linked to both the number of cases and the severity of acne. When you use comedogenic or improper skin care products, they can block your pores and make acne worse. This shows that more people need to know how to choose goods that are non-comedogenic and dermatologically tested, especially those who use cosmetics a lot.

The multivariate logistic regression analysis added to these results by showing that worry, poor diet, lack of sleep, and cosmetic use are all independent predictors of acne vulgaris. These results show that acne isn't just a biological problem, but is also affected by lifestyle choices that can be changed. So, treatments that focus on these factors might help lower the number of people who get acne and make it less severe. It's interesting that there was no significant link found between the prevalence of acne and age or school year. This means that acne affects students in this age range in a pretty uniform way. There was a weak negative relationship

between physical exercise and acne, but it wasn't strong enough to be statistically significant. There could be different ways people exercise or other things that skewed the results that weren't taken into account in this study [13].

This work has some good points, like clinical assessment and multivariate analysis, but it also has some problems. It is hard to find direct links between risk factors and pimples because the study is cross-sectional. There is also a chance of selection bias when convenience sampling is used, and recall bias can happen when people describe their own lifestyle factors. Also, the study was only done at one school, which means that the results may not be applicable to other groups of people.

Conclusion

The results show that acne isn't just a physical problem; it's also affected by things like stress, food, sleep, and cosmetic use that can be changed. Stress was the most important indicator, followed by bad eating habits and not getting enough sleep. This shows that acne is caused by more than one thing.

These results show how important it is to take a more complete approach to treating acne, one that includes changes to your lifestyle as well as traditional drug-based treatments. A big part of lowering the number of young adults with acne could be educational programs that teach them how to deal with stress, eat well, get enough sleep, and take care of their face properly.

The study also shows how important it is to make students more aware of how their daily habits can affect their skin health. To find out what causes what and how to make targeted, evidence-based prevention and management plans, future research should focus on longitudinal and multi-center studies.

References

1. Alsaadoon, Noor Sulaiman Jalood, Asmaa Mohammed Al-Refaie, and Aml Youssif Habashy. "Acne vulgaris in adolescents: A comprehensive review." *Benha Journal of Applied Sciences* 9, no. 12 (2024): 5-13.
2. Guguluş, Dumitrița Lenuța, Dan Vătă, Ioana Adriana Popescu, Adriana Ionela Pătrașcu, Ioana Alina Halip, Mădălina Mocanu, and Laura Gheucă Solovăstru. "The epidemiology of acne in the current era: trends and clinical implications." *Cosmetics* 12, no. 3 (2025): 106.
3. Akl, Essam Mohamed, Samah Ezzat Ibrahim, Nehad Ahmed Fouad, and Ebtessam Ahmed Mowafy. "Etiopathogenesis of acne vulgaris." *Benha Journal of Applied Sciences* 9, no. 7 (2024): 7-10.

4. Olceroglu, Bilge, Gamze Balci, and Ahmet Kati. "The Skin Microbiome and Its Implications for Maintaining Skin Health." In *Phytochemical Potentials for Dermatological Applications*, pp. 76-103. CRC Press, 2025.
5. Schwingshackl, Lukas, Lisa Patricia Hobl, and Georg Hoffmann. "Effects of low glycaemic index/low glycaemic load vs. high glycaemic index/high glycaemic load diets on overweight/obesity and associated risk factors in children and adolescents: a systematic review and meta-analysis." *Nutrition journal* 14, no. 1 (2015): 87.
6. Jović, Anamaria, Branka Marinović, Krešimir Kostović, Romana Čević, Aleksandra Basta-Juzbašić, and Zrinka Bukvić Mokos. "The impact of psychological stress on acne." *Acta Dermatovenerologica Croatica* 25, no. 2 (2017): 133-133.
7. Prakesh, Utter, and Mahmood Abbas. "The Rising Incidence of Acne Vulgaris in Adolescents: Lifestyle Factors and Preventive Strategies in Bangladesh." *Scientific Journal of Dermatology and Venereology* 3, no. 1 (2025): 54-66.
8. Dunn, Lauren K., Jenna L. O'Neill, and Steven R. Feldman. "Acne in adolescents: Quality of life, self-esteem, mood and psychological disorders." *Dermatology online journal* 17, no. 1 (2011).
9. Mistry, Durjoy, Mehjabin Hossain, B. S. Diba, J. D. Plabon, M. A. Shams, M. T. Rahman, and A. S. M. Bakibillah. "A Multisectoral Study of Mpox Epidemiology, Resistance Surveillance, and Policy Gaps: Toward a One Health Framework." *Clinical Reviews in Allergy & Immunology* 68, no. 1 (2025): 108.
10. Currie, Dawn, and Sara Wiesenber. "Promoting women's health-seeking behavior: research and the empowerment of women." *Health care for women international* 24, no. 10 (2003): 880-899.
11. Tripathy, Krisha, Ajay S. Dulai, Mildred Min, and Raja K. Sivamani. "Insulin Resistance, Metabolic Syndrome, and Inflammatory Skin Disease." *Journal of Clinical Medicine* 15, no. 1 (2026): 330.
12. Menge, Jordan. "How to Protect Your Skin from Stress & Sleep Disruption."
13. Kunin, Audrey. *The DERMAdoctor skininstruction manual: The smart guide to healthy, beautiful skin and looking good at any Age*. Simon and Schuster, 2007.