

Biomedical and Biotechnological Sciences

Research Article

Toxicological Assessment of Supplemental *Zanthoxylum acanthopodium* essential oil in Rabbits: Evidence from Haemato-biochemical and Histopathological Indices

Alagbe J.O^{1*}, Anaso, E.U.², Shittu, M.D.³, Anuore, D.N⁴, Emiola, I.A⁵

¹Adjunct Lecturer, Department of Biochemistry, Gandhi College of Agriculture, Rajasthan, India

¹Assistant Professor, Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India

²Department of Animal Science, Federal University of Agriculture, Mubi Adamawa State, Nigeria

³Department of Animal Production and Health, Ladoke Akintola University of Technology, Ogbomosho, Nigeria

⁴Department of Animal Science, University of Abuja, Gwagwalada, Nigeria

⁵Department of Animal Nutrition and Biotechnology, Ladoke Akintola University of Technology, Ogbomosho, Nigeria

***Corresponding Author:** Alagbe J.O, Adjunct Lecturer, Department of Biochemistry, Gandhi College of Agriculture, Rajasthan, India.

Received Date: 12 January 2026; **Accepted Date:** 02 February 2026; **Published Date:** 11 February 2026

Copyright: © 2026 Alagbe J.O, this is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

This study investigated the toxicological assessment of supplemental *Zanthoxylum acanthopodium* essential oil in rabbits: evidence from haemato-biochemical and histopathological Indices. Thirty two clinically healthy weaner male rabbits of about 7 to 8 weeks with an average initial weight live weight of 710 ± 0.68 kg were stratified by body weight and randomly allocated to one of four experimental groups (n = 8). The experimental diet consisted of a basal (control) group (0 mL/kg) and three levels of *Zanthoxylum acanthopodium* essential oil inclusion: 0.5 mL, 1.0 mL and 1.5 mL per kg DM feed daily. Feed and water was made available at all times and a completely randomized design was adopted throughout the 14 weeks experimental period during which rabbits were maintained under standard management conditions. The results suggested that *Zanthoxylum acanthopodium* essential oil influenced ($p < 0.05$) several haematological parameters. Specifically, haemoglobin, red blood cell, mean corpuscular volume, mean corpuscular haemoglobin and mean corpuscular haemoglobin concentrations. Conversely, pack cell volume, white blood cell, lymphocytes, monocytes, neutrophils, basophils and eosinophils levels were not affected by the treatment. In serum analysis, total protein, albumin, globulin, glucose, creatinine, sodium, potassium, alkaline phosphatase, alanine amino transferase and aspartate amino transferase remained stable across all groups ($p > 0.05$); however, a significant effect was observed on cholesterol, calcium, phosphorus and magnesium levels. Liver histology was not affected by *Zanthoxylum acanthopodium* essential oil across all treatments showing no signs of toxicity or degeneration. Furthermore, all blood values remained within the recommended physiological ranges for healthy rabbits. The study concludes that *Zanthoxylum acanthopodium* essential oil can be safely incorporated into rabbit diets up to 1.5 mL as it supports normal physiological function and maintains liver integrity.

Keywords: *Zanthoxylum acanthopodium* oil, Rabbits, Haematology, Serum biochemistry, Hepatic histology, Cholesterol.

Introduction

Globally, rabbit production is experiencing significant growth particularly in developing countries due to the high prolificacy, rapid growth rate and superior nutritional quality of rabbit

meat [1]. To maximize these production benefits, nutritionist have shifted focus towards the use of phyto-genic feed additives – natural plant extracts such as essential oils to improve

animal sustainability, health and performance [1]. This transition is largely driven by worldwide trend of reducing synthetic antibiotics use in animals due to concerns over antimicrobial resistance and toxic residues in animal products [3,4].

Essential oils are volatile, aromatic compounds extracted from herbal or medicinal plants that possess well documented anti-inflammatory, antioxidant, anti-helminthic, cytotoxic, hypolipidemic, antidiarrheal, antifungal, immune-modulatory, gastro-protective, dermato-protective, cardio-protective, antimicrobial, hepato-protective and antidiabetic properties [1,5]. While their potential to improve growth performance and feed efficiency is widely recognized, it is imperative to establish their safety profile regarding the internal physiology of the animal [6,7]. Haematological and serum biochemical parameters are important indicators of the nutritional, physiological and pathological status of animals [8]. Deviations in these parameters often serve as a systemic toxicity or metabolic stress [9,10]. Furthermore, because the liver is the primary organ responsible for detoxification of ingested substances, its histological is a critical standard for evaluating the safety of any dietary inclusion [11,12].

Despite the promising benefits of essential oils, there is need to determine the optimal inclusion levels that can be tolerated without compromising the animal's health. Previous research by [13,14] have shown that the supplementation of clove and eugenol oil at low doses below 500 mg/kg in the diet of rabbits revealed beneficial effects, high doses above 500 mg/kg can lead to liver damage. Therefore, some caution should be exercised in the use essential oils as natural feed additives in rabbit diets. Recent studies on phytochemicals, although very few, have been carried using different plant extracts on animal's performance. Whereas essential oils from rosemary, thyme, clove, ginger, turmeric, amongst others have been investigated [14,15,9]. However, none has investigated the toxicological assessment of supplemental Zanthoxylum acanthopodium essential oil in rabbits: evidence from haemato-biochemical and histopathological Indices.

Materials and Methods

Experimental location, Ethical Approval and Animal care

The experiment was carried out at the Rabbit unit of the Gandhi College of Agricultural Teaching and Research Farm, Rajasthan, India. The study site lies between latitude 11° 25'N and 16° 00'E and longitude 4° 00'N and 9° 07'E. The mean annual rainfall and temperature range from 400 to 1000 mm and 21.83 to 32.57°C, respectively. Relative humidity is about 70 % during the raining season and 45 % during the dry season. The regulations relating to animal care and the use of animals in this experiment were approved by the Research and Ethics Committee within the Department of Animal Nutrition and Biochemistry, Gandhi College of Agriculture, Rajasthan India.

Sanitation commenced two weeks before the purchase

of experimental animals with thorough cleaning and disinfection of pens and other equipment's with disinfectant (Morigad). Feeders were properly washed and nipple drinkers was flushed with antiseptic (Aquaclean). Thirty two clinically healthy weaner male rabbits of about 7 to 8 weeks sourced from a reputable source in Rajasthan India. They have an average initial weight live weight of 710 ± 0.68 kg were stratified by body weight, such that the rabbits in each treatment group had similar average initial body weight, and randomly allocated to one of four experimental groups ($n = 8$). On arrival, rabbits were placed on a two weeks adjustment period and dewormed with Ivermectin Plus (Needan Pharmaceuticals, Gujarat India) at the dosage of 1 mL to 1 kg BW. Animals were housed individually in an all wired battery cage equipped with galvanized feeders for each animal and nipples for supplying fresh water. Basal diet was compounded according to the nutritional standard described by [16]. A completely randomized experimental design was adopted. The dietary treatments were: (A) Basal diet without a supplement (control); (B) Basal diet with Zanthoxylum acanthopodium essential oil supplement (per kg DM feed daily) at 0.5 mL; (C) Basal diet with Zanthoxylum acanthopodium essential oil supplement (per kg DM feed daily) at 1.0 mL; (D) Basal diet with Zanthoxylum acanthopodium essential oil supplement (per kg DM feed daily) at 1.5 mL. Feed was given trice daily ad libitum at 07:30 h, 12:00 h and 17:30 h. The feeding trial lasted for 14 weeks including two weeks acclimatization period. Before the first feeding, each rabbits received a 100 g feed with dose of essential oil to guarantee full consumption. Proximate analysis of basal diet was ascertained according to [17] procedure.

Zanthoxylum acanthopodium leaf collection, extraction and GC-MS analysis

Fresh leaves of Zanthoxylum acanthopodium were harvested from different trees at Gandhi College of Agriculture, Rajasthan India. It was sent to the department of Biological Sciences in the same institution where it was identified by a certified taxonomist (Dr. Ram Vinod) and registered under voucher number HY/004/2023. Leaves were shade dried for 10 days and pulverized using mechanical grinder. Extraction of oil was done by hydrodistillation with an H-shaped Clevenger-type apparatus according to the procedures modified by [1,18]. Briefly, 300 g of the pulverized Zanthoxylum acanthopodium was added to 1200 mL of water heated in a glass flask at 60 °C for 20 minutes, steam passes via the condenser and collected in a beaker. The oil was collected by decantation and then dried over column of anhydrous sodium sulfate before it was introduced into glass bottles and stored in a refrigerator at 4 °C.

GC/MS analysis of Zanthoxylum acanthopodium essential oil was done using Claudius 5006 GC-MS Auto Sampler (China) equipped with two silica capillary columns, interfaced with a quadrupole detector (single quadrupole acquisition

Method-MS parameters report), source temperature 230°C, Quadrupole temperature 150 °C; the temperature program was 60 °C for 2 min, 60-240 °C at 3 °C/min, then kept at 240 °C during 8 min; injector temperature, 240 °C. The mass spectrometry transfer line temperature, 250 °C; carrier gas, helium at a flow rate of 0.7 ml/min; injection type, split, 20:1; ionization voltage, 70 eV; electron multiplier 1000 eV; scan range 33-400 amu; scan rate, 1.56 scan/s.

Identification of components Interpretation on mass spectrum GC-MS was conducted using the database of National Institute Standard and Technology (2001) having more than 62,000 patterns. The spectrum of the unknown component was compared with the spectrum of the known components stored in the NIST library.

Histology Analysis

On the last day of the experiment, 3 animals were randomly selected for histological study. The animals were slaughtered and their liver was removed, fixed in 10 % neutral-buffered formalin, dehydrated in 70% alcohol, cleared with methyl benzoate, and embedded in paraffin wax. Sections of 5µm were cut and stained on glass slides with hematoxylin and eosin stain for light microscopic examination. Stained sections were examined by light microscope and photographed using a digital camera.

Collection of Blood for Analysis

On the last day of the trial, 2 sets of blood were collected very early in the morning from four randomly selected rabbits in each treatment through vein using a 5 mL syringe. 5 ml blood sample was collected into labelled sterile bottles containing disodium salt of ethylene diamine tetra-acetic acid (EDTA) as anticoagulant for the determination of haematological parameters. Blood samples for serum analysis were collected into anticoagulant free bottles. All samples collected were placed in an ice pack and sent to the laboratory for further analysis. Blood for haematology was analyzed using BC-5390 Auto Hematology Analyzer (China). The Kit utilizes Impedance method for pack cell volume, mean corpuscular haemoglobin concentration, red blood cell, mean corpuscular determination, Cyanide free reagent for hemoglobin test and Flow Cytometry (FCM) + Laser scatter + Chemical dye method for white blood cell and its differentials analysis. Blood for serum was analyzed using Sysmex Auto Biochemical analyzer – HD -3040 Series. The sample rotor valve measures the required volume of sample precisely to provide accurate and reliable results for all serum biochemical parameters.

Data Analysis

Data obtained were subjected to analysis of variance (ANOVA) for a complete randomized design using Statistical Package for the Social Sciences (SPSS version 25). When the ANOVA was significant, means were separated using Duncan's multiple range test at the level of $P \leq 0.05$.

Results

Ingredients	(kg of DM)
Corn	35.00
Wheat bran	20.00
Palm kernel meal	16.29
Soybean meal	20.06
Limestone	2.50
Bone meal	5.00
Lysine	0.20
Methionine	0.20
Min/Vitamin Premix	0.25
Salt	0.35
Toxin binder	0.15
Total	100.0
Chemical composition	
Dry matter	91.25
Organic matter	90.14
Crude protein	16.21
Crude fibre	13.87
Ash	9.86
Energy (Kcal/kg)	2891.5

Each 2.5 kg consists of: Vit A 12000, 000 IU; Vit D3, 2000, 000 IU; Vit. E. 10g; Vit k3 2 g; Vit B1, 1000 mg ; Vit B2, 49g ; Vit B6, 105 g; Vit B12, 10 mg; Pantothenic acid, 10 g; Niacin, 20 g , Folic acid , 1000 mg ; Biotin, 50 g; Choline Chloride, 500 mg, Fe, 30 g; Mn, 40 sg; Cu, 3 g; Co, 200 mg; Si, 100 mg and Zn , 45 g

Table 1: Ingredient and chemical composition of the basal diet (kg of DM)

Compounds	Retention time (minutes)	% Area
9,17-Octadecadienal	6.113	10.77
alpha-copaene	7.008	6.55
γ-Terpinene	7.072	8.18
p-Cymene	8.009	5.72
Linalool	10.76	12.86
α-Humulene	10.95	3.89
Terpinene-4-ol	10.99	4.43
Limonene	11.85	16.85
β-Caryophyllene	12.10	14.37
β-Citronella	12.94	5.14
cis-Vaccenic acid	13.88	2.09
9-Eicosenoic acid	14.12	3.11

Table 2: Dominant bioactive compounds in Zanthoxylum acantho-

podium oil

Dominant bioactive compounds in *Zanthoxylum acanthopodium* oil is presented in Table 2. Twelve major bioactive compounds were identified including, Limonene (16.85 %), β -Caryophyllene (14.37 %), Linalool (12.86 %), 9, 17-Octadecadienal (10.77 %), γ -Terpinene (8.18 %), α -copaene (6.55 %), p-Cymene (5.72 %), β -Citronella (5.14 %), Terpinene-4-ol (4.43 %), α -Humulene (3.89 %) amongst others.

Haematological parameters of weaner rabbits fed diet supplemented with *Zanthoxylum acanthopodium* oil is presented in Table 3. Haemoglobin, red blood cell, mean corpuscular volume, mean corpuscular haemoglobin and mean corpuscular haemoglobin concentration were higher ($p < 0.05$) for diet B (DB), diet C (DC) and diet D (DD) than for diet A (control). Pack cell volume, white blood cell, lymphocytes, monocytes, basophils and eosinophil values were similar ($p > 0.05$) among the diets.

Components	DA	DB	DC	DD	SEM	P-value
Packed cell volume (%)	32.92	33.56	33.97	34.05	0.85	0.275
Hemoglobin (g/dL)	9.66b	11.72a	11.86a	11.91a	2.21	0.001
Red blood cells (106/ L)	13.87b	14.25a	14.37a	14.40a	0.56	0.899
MCHC (%)	28.17b	34.86a	34.98a	35.07a	0.71	0.002
MCH (pg)	12.31b	18.74a	18.82a	18.93a	3.08	0.007
Mean corpuscular volume(fL)	48.87b	57.11a	58.07a	58.91a	6.65	0.933
White blood cells (109/L)	13.55	13.65	13.93	14.04	2.76	0.406
Lymphocytes (%)	60.34	66.85	67.12	67.24	7.07	0.587
Monocytes (%)	2.95	2.76	2.80	2.86	0.41	0.665
Neutrophils (%)	45.17	47.08	47.11	47.23	5.94	0.872
Basophils (%)	0.07	0.06	0.06	0.06	0.02	0.961
Eosinophils (%)	4.00	4.03	4.11	4.15	1.1	0.751

Table 3: Haematological parameters of weaner rabbits fed diet supplemented with *Zanthoxylum acanthopodium* oil

Parameters	DA	DB	DC	DD	SEM	P-value
Total protein (g/L)	69.12	71.33	71.47	71.81	7.86	0.61
Albumin (g/L)	33.51	34.05	34.12	34.15	4.31	0.825
Globulin (g/L)	35.61	37.28	37.35	37.66	3.46	0.512
Albumin:globulin ratio	0.94	0.91	0.91	0.90	0.10	0.801
Glucose (mg/dL)	70.76	70.75	70.72	70.73	0.03	0.993
Cholesterol (mg/dL)	117.8a	89.34b	89.11b	88.75b	2.08	0.882
Urea (mg/dL)	4.16	4.05	4.11	4.09	1.05	0.772
Creatinine (mg/dL)	2.36	2.57	2.67	2.77	0.11	0.761

Means within a row with different letters and significantly different ($p < 0.05$); SEM Standard error; T1: Experimental diet without oil (control); T2: experimental diet + 2.5 g Adriamycin® /kg diet; T3: experimental diet + 2.0 mL *Clinopodium brownei* oil/kg diet; T4: experimental diet + 2.0 mL *Clinopodium brownei* oil/kg diet

Table 4: Table 5: Serum biochemical indices of weaner rabbits fed diet supplemented with *Zanthoxylum acanthopodium* oil

Except for cholesterol concentration which was influenced ($p < 0.05$) by the treatment, value was higher in DA than in other treatments. Total protein, albumin, globulin, albumin: globulin ratio, glucose, creatinine and urea concentration showed no ($p > 0.05$) significant difference (Table 4). Serum ions and enzymes of weaner rabbit fed diet supplemented

with *Zanthoxylum acanthopodium* oil is presented in Table 5. Except for calcium, phosphorus, magnesium which were influenced ($p < 0.05$) by the treatments, other parameters (sodium, potassium, alkaline phosphatase, alanine amino transferase, aspartate amino transferase) were not affected ($p > 0.05$) by the treatments.

Items	DA	DB	DC	DD	SEM	P-value
Calcium (mmol/L)	2.24b	3.56a	4.72a	4.81a	0.02	0.175
Phosphorus (mmol/dL)	4.77b	6.96a	7.02a	7.05a	0.31	0.811
Magnesium (mmol/L)	2.05b	3.81a	3.98a	4.01a	0.12	0.183
Sodium (mmol/L)	149.6	150.8	151.7	153.1	4.47	0.900
Potassium (mmol/L)	3.67	3.83	3.91	3.96	0.35	0.586
Alkaline phosphatase (IU/L)	29.11	28.59	28.07	28.08	0.12	0.834
Alanine amino transferase (IU/L)	16.57	15.95	15.08	15.16	0.69	0.115
Aspartate amino transferase (IU/L)	58.55	51.03	51.25	51.20	4.04	0.806

Table 5: Serum ions and enzymes of weaner rabbit fed diet supplemented with *Zanthoxylum acanthopodium* oil

Discussions

Bioactive compounds in *Zanthoxylum acanthopodium* oil has numerous medicinal properties including, anti-inflammatory, dermato-protective, gastro-protective, immuno-stimulatory, cardio-protective, antioxidant, antimicrobial, hepato-protective, cytotoxic, anti-helminthic, antidiabetic, antidiarrheal, hypolipidemic, amongst others [19,20]. These phyto-compounds helps to improve the overall health of rabbits because they are eco-friendly, contains non-toxic compounds and generally regarded as safe [14,21]. Additionally, it is well documented that dietary supplementation of essential oils in the diet of rabbits modulates the intestinal flora, immune activity and prevent the incidence of antimicrobial resistance [22,23].

All the haematological indices obtained in this study were within the normal physiological ranges for healthy rabbits [24]. The red blood cell, haemoglobin and pack cell volume were within the normal range 8.99 – 16.00 (106/ L), 7.00 – 14.00 g/dL and 29.00 – 39.00 % recorded by [14]. When red blood cell, haemoglobin and pack cell volume are all within the normal range, it implies the absence of anaemia, oxygen sufficiency in the tissues as well as efficient hydration [25,20]. Mean corpuscular volume, mean corpuscular haemoglobin and mean corpuscular haemoglobin concentration were within 41.00 – 67.00 fl, 17.00 – 24.00 pg and 30.00 – 36.00 % cited by [26]. These outcome indicates the absence of chronic iron deficiency, inflammation as well as folate deficiency in the blood [27]. White blood cell protects the body cell from damage and helps to maintain proper immune function [28]. As earlier discussed, essential oil possess antioxidant properties thus inhibiting the efficacy of pathogenic organisms in the gut as well as neutralizing the activities of free radicals which are the cause of most chronic diseases [29]. However, their value was within 12.00 – 25.00 (109/L) reported by [30]. This outcome further to reveal that the animals used for this current study were healthy [31]. Lymphocytes, monocytes, neutrophils, basophils and eosinophil counts were within the normal ranges, 40.00 – 78.00 %, 1.00 – 3.00 %, 30.00 –

50.00 %, 0.01 – 1.00 % and 0.1 – 5.00 % reported by [32]. This result suggests a stability in the immune system as well as absence of tissue inflammation [33]. The presence of limonene, β -caryophyllene, linalool and other bioactive compounds in *Zanthoxylum acanthopodium* oil are probably the reason for the strong immune capacity in rabbits. [33]

All the serum parameters were within the normal physiological ranges for healthy rabbits.[6] This results indicates a balanced metabolic state in rabbits. As previously noted, the bioactive compounds in *Zanthoxylum acanthopodium* oil have antimicrobial activities against some bacteria's ensuring that the animals internal organs are functioning properly [30]. Total protein, albumin and globulin values were within the reference range 40.00 – 85.00 g/L, 28.00 – 40.00 g/dL and 20.00 – 50.00 g/L cited by [34]. Normal albumin and globulin levels in rabbits suggests efficient protein synthesis by the level which translates to a better performance and weight gain [35]. Normal globulin range also indicates the absence of chronic inflammatory diseases [36]. This result concur with earlier studies [37]. A normal glucose level 70.00 – 155 mg/dL is a clear sign that the rabbits were not severely stressed due to malnutrition and other management practices [2]. Cholesterol level recorded in this study was within 97.00 – 180.5 mg/dL recorded by [37]. Recently, Alagbe and Hernandez (2025) observed that essential oils have the ability to reduce the concentrations of saturated fatty acid and prevent cardio-vascular diseases when supplemented in the diet of animals. The lowered cholesterol levels among animals which received *Zanthoxylum acanthopodium* oil may be due to the presence of bioactive compounds especially cis-Vaccenic acid and 9-Eicosenoic acid which have been previously associated with hypolipidemic properties [37]. Creatinine and urea concentrations are most reliable marker for kidney degeneration [31]. However, urea and creatinine ranges of 4.09 – 4.11 mg/dL and 2.36 – 2.77 mg/dL recorded in this study was within 2.50 – 5.60 mg/dL and 2.00 – 4.50 mg/dL referenced by [30]. This outcome suggests the ab-

sence of chronic renal disease in rabbits [38]. The normal range of serum calcium (1.88 – 4.33 mmol/L; [37], phosphorus (4.00 – 8.50 mmol/L; [39], magnesium (2.00 – 4.10 mmol/L; [30], sodium (100 – 210 mmol/L; [40] and potassium (2.10 – 3.80 mmol/L; [41] for all treatments suggests that *Zanthoxylum acanthopodium* oil supplementation in the diet of rabbits did not interfere the absorption of minerals in the tissues. This outcome is parallel with the reports of [24,42] who recorded that the inclusion of phyto-

genics in the diet of growing rabbits did not affect the absorption and availability of nutrients in their system. Although alkaline phosphatase (ALP), alanine amino transferase (AST) and aspartate amino transferase (ALT) were not influenced ($p>0.05$) by the treatment diets, values were within the range of 18.00 – 30.00 IU/L, 12.00 – 25.00 IU/L and 45.00 – 70.00 IU/L reported for health growing rabbits [43]. ALT, ALP and AST normal levels suggests no hepato-cellular damage (Ei-




Plate 1: Basal diet without essential oil supplement (control)




Plate 2: Basal supplemented with supplemented with *Zanthoxylum acanthopodium* oil at 0.5 ml/kg diet




Plate 3: Basal supplemented with supplemented with *Zanthoxylum acanthopodium* oil at 1.0 ml/kg diet




Plate 4: Basal supplemented with supplemented with *Zanthoxylum acanthopodium* oil at 1.5 ml/kg diet

Hepatic histology of weaner rabbits fed diet supplemented with *Zanthoxylum acanthopodium* oil is presented in plate 1-4. Plate 1 (control) without oil supplement, plate 2, 3 and 4 contained basal diet supplemented with *Zanthoxylum acanthopodium* oil at 0.5 mL, 1.0 mL and 1.5 mL/kg diet respec-

tively. There was no significant ($p > 0.05$) in their histological examinations, all slides show normal hepatic architecture without congestions, there was normal or clear radial arrangement of hepatic cords around the central vein without any signs of inflammation or fatty infiltration and well-formed

sinusoids. The outcome of this result suggests that dietary supplementation of *Zanthoxylum acanthopodium* oil up to 1.5 ml/ kg diet was optimum for the animals and had no toxic effect on their liver histology. This result is in line with the observations of [44] who noted a normal central vein and hepatic cords without inflammation with the dietary supplementation of ginger, clove, red pepper and black pepper in growing rabbits. Additionally, it is well documented that supplementing diets of animals with essential oils pose no vacuolar degeneration [45, 25] noted that the administration of mixtures of essential oils ensures that the hepatocyte plasma membrane and portal lobules were well organized. Aside toxicity, environmental stress have also been linked to one of the causes of minor shift in hepatic histology [46]. However, bioactive compounds in *Zanthoxylum acanthopodium* oil have proven to possess protective or therapeutic effect when growing rabbits are exposed to toxins and environmental stressors (Volex et al., 2018).

Conclusion

The outcome of this study demonstrates that dietary supplementation of *Zanthoxylum acanthopodium* oil up to level of 1.5 mL/kg is physiologically safe and well tolerated by growing rabbits while the oil significantly influenced some blood parameters it did not trigger an adverse immune response or systemic inflammation. Thus this study warrant further investigation.

References

1. Sharma, Singh, Alagbe Olujimi John, Liu Xing, Sharma Ram, and Kumar Amita. "Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroemetry." *International Journal of Agriculture and Animal Production* 2, no. 6 (2022): 18-27.
2. Alagbe, John O and Oluwafemi, R.A (2019). Hematology and Serum Biochemical Indices of Growing Rabbits Fed Diet Supplemented with Different Level of *Indigofera Zollingeriana* Leaf Meal. *Progress in Chemical and Biochemical Research* 2019, 2(4), 170-177.
3. Alagbe, John Olujimi. "Impact of dietary supplementation of *Rhamnus prinoides* leaf extract on the growth performance, nutrient retention and intestinal microbial count of "japanese quails"." *Brazilian Journal of Science* 3, no. 5 (2024): 40-50.
4. Ojedian, Taiwo Kayode, I. A. Emiola, V. Durojave, and John Olujimi Alagbe. "Proximate, vitamin and GC-MS profiling of *Kigelia africana* fruit powder." *Cerrado: Agricultural and Biological Research* 1, no. 1 (2024): 13-20.
5. Alagbe, John Olujimi. "Clerodendron splendens leaf extract supplementation in weaner rabbits: impact on growth performance, haematology and intestinal microbial population." *Cerrado: Agricultural and Biological Research* 1, no. 1 (2024): 21-31.
6. Alagbe, J. O., E. U. Anaso, H. Zubairu, and D. N. Anorue. "Lep-tadenia Hastate and Cedrus Brevifolia Oil: Effect on Growth Performance, Caecal Microbial Population, Fermentation and Haematological Indices of Weaner Rabbits." *International Journal of Advanced Technology and Social Sciences* 3, no. 9 (2025): 1213-1228.
7. Hernandez, M and Alagbe, J.O. (2025a). Influence of Abrus procatorious crude oil supplementation on the growth performance, Nutrient digestibility, Ruminant fermentation and Microbial population of Malabari Bucks. *International Journal of Global Sustainable Research*, 3(7): 527-538.
8. Morris, Hernández, and Alagbe John Olujimi. "INFLUENCE OF *Odontonema Strictum* OIL ON THE GROWTH PERFORMANCE AND RUMINAL FERMENTATION OF BARBARI BUCKS." *Research in: Agricultural & Veterinary Sciences* 9, no. 2 (2025).
9. Yang, C. J., I. Y. Yang, D. H. Oh, I. H. Bae, S. Goo Cho, I. G. Kong, D. Uganbayar, I. S. Nou, and K. S. Choi. "Effect of green tea by-product on performance and body composition in broiler chicks." *Asian-Australasian Journal of Animal Sciences* 16, no. 6 (2003): 867-872.
10. Alagbe, John Olujimi. "Effect of coconut shell extract on the growth performance and some haemato-biochemical parameters of broiler chicken." *Brazilian Journal of Science* 3, no. 6 (2024): 82-95.
11. El-Gogary, M. R., E. A. El-Said, and A. M. Mansour. "Physiological and immunological effects of rosemary essential oil in growing rabbit diets." *J. Agric. Sci* 10 (2018): 485-491.
12. Anna C., Renata klebaniuk, Eugeniusz Ryszard G., Wioletta Samolińska, Katarzyna Ognik (2017). Polish crossbred pigs' blood haematological parameters depending on their age and physiological state. *Annals of Warsaw University of Life Sciences – SGGW Animal Science* 56 (2): 185–195.
13. Bassiony, Samar SM, M. M. Elhindawy, A. E. Attia, and I. E. Ismail. "Effect of some bioactive components of essential oils on growing rabbits performance." *Zagazig Journal of Agricultural Research* 42, no. 5 (2015).
14. Abdelnour, Sameh, Mahmoud Alagawany, Mohamed E. Abd El-Hack, Asmaa M. Sheiha, Islam M. Saadeldin, and Ayman A. Swelum. "Growth, carcass traits, blood hematology, serum metabolites, immunity, and oxidative indices of growing rabbits fed diets supplemented with red or black pepper oils." *Animals* 8, no. 10 (2018): 168.
15. Corduk, M., S. Sarica, and G. F. Yarim. "Effects of oregano or red pepper essential oil supplementation to diets for broiler chicks with delayed feeding after hatching. 1. Performance and microbial population." *Journal of Applied Poultry Research* 22, no. 4 (2013): 738-749.
16. National Research Council (NRC). *Nutrient Requirements of Rabbits*, 2nd ed.; *National Academy Press: Washington, DC, USA*, 1977.

17. AOAC. 2003. Association of Official Analytical Chemists Official Methods of Analysis 15th ed. Published the AOAC, Washington, DC, USA
18. Anaso, Emmanuel U., Olurotimi A. Olafadehan, Ijeoma C. Chibuogwu, and John O. Alagbe. "Seminal morphology and organ morphometrics of rabbit bucks fed Piliostigma thonningii essential oil supplemented diet." *Science Letters* 12, no. 2 (2024): 70-75.
19. Ojediran, Taiwo Kayode, Olujimi John Alagbe, Durojaye Victor, and Emiola Adewale. "Analysis of *Kigelia africana* (Lam.) Benth. fruit powder's antioxidant and phytochemical properties." *Brazilian Journal of Science* 3, no. 7 (2024): 38-49.
20. Alagbe, John Olujimi. "Effect on performance, serum biochemistry and haematological components of feeding "japanese quails" phytogenic feed additions comprising *Megaphrynium macrostachyum* leaves." *Brazilian Journal of Science* 3, no. 5 (2024): 51-64.
21. Alagbe, J. O. "Characterization Of Bioactive Compounds in *Luffa Aegyptiaca* Leaf Ethanolic Extracts Using Gas Chromatography And Mass Spectrometer: Characterization Of Bioactive Compounds in *Luffa Aegyptiaca* Leaf Ethanolic Extracts Using Gas Chromatography And Mass Spectrometer." *Food Science & Applied Microbiology Reports* 1, no. 2 (2022): 21-28.
22. Nouzarian, R., S. A. Tabeidian, M. Toghyani, G. Ghalamkari, and M. Toghyani. "Effect of turmeric powder on performance, carcass traits, humoral immune responses, and serum metabolites in broiler chickens." *J. Anim. Feed Sci* 20, no. 3 (2011): 389-400.
23. Shittu, M. D., and J. O. Alagbe. "Phyto-nutritional profiles of broom weed (*Sida acuta*) leaf extract." *Annals of Clinical Toxicology* 3, no. 2 (2020).
24. Shittu, M. D., O. J. Alagbe, O. Alaba, E. O. Okanlawon, F. A. Adelokun, P. O. Emmanuel, and D. O. Adejumo. "Effect of ginger, garlic and negro pepper on gut microbes, gut histomorphometry and pathological assessment of selected organs of broiler chickens." *ADAN JOURNAL OF AGRICULTURE* 5, no. 1 (2024).
25. Ashour, E., M. Alagawany, F. Reda, and M. Abd El-Hack. "Effect of supplementation of *Yucca schidigera* extract to Growing rabbit diets on growth performance, carcass characteristics, serum biochemistry and liver oxidative status." *Asian J. Anim. Vet. Adv* 9 (2014): 732-742.
26. Dalle Zotte, Antonella, C. Celia, and Zs Szendrő. "Herbs and spices inclusion as feedstuff or additive in growing rabbit diets and as additive in rabbit meat: A review." *Livestock science* 189 (2016): 82-90.
27. Bakeer, Manal Ramadan. "Focus on the effect of dietary pumpkin (*Cucurbita moschata*) seed oil supplementation on productive performance of growing rabbits." *Journal of Applied Veterinary Sciences* 6, no. 2 (2021): 22-26.
28. Oloruntola, Olugbenga D., Simeon O. Ayodele, Samuel Adebowale Adeyeye, Deborah Adebukola Oloruntola, Clement O. Osowe, and Oluwagbemiga Samuel Fasuhami. "Broiler Chickens' Growth, Haematological Indices, Guts Microbiota, Carcass and Meat Analysis in Response to Dietary Supplementation with *Anacardium occidentale* Leaf Powder and A Mix of Prebiotic, Probiotic and Acidifier." *Journal of Poultry Research* 19, no. 2 (2022): 52-59.
29. Anorue, Daniel Nnadozie, Friday Ubong, and Alagbe Olujimi John. "Investigating the effects of pawpaw (*Carica papaya*) essential oil dietary supplementation on the growth performance and carcass characteristics of broilers." *Research in: Agricultural and Veterinary Sciences* 7, no. 3 (2023): 164-174.
30. Bacova, Kristina, Karin Zitterl-Eglseer, Lubica Chrastinova, Andrea Laukova, Michaela Madarova, Sona Gancarcikova, Drahomira Sopkova, Zuzana Andrejckakova, and Iveta Placha. "Effect of thymol addition and withdrawal on some blood parameters, antioxidative defence system and fatty acid profile in rabbit muscle." *Animals* 10, no. 8 (2020): 1248.
31. Liu, Yang, Yang Li, Jiaying Niu, Hua Liu, Ning Jiao, Libo Huang, Shuzhen Jiang, Lei Yan, and Weiren Yang. "Effects of dietary *Macleaya cordata* extract containing isoquinoline alkaloids supplementation as an alternative to antibiotics in the diets on growth performance and liver health of broiler chickens." *Frontiers in Veterinary Science* 9 (2022): 950174.
32. Alagawany, Mahmoud, Mohamed S. Elewa, Diaa E. Abou-Kassem, Tamer Ahmed Ismail, Ayman S. Salah, Mahmoud Madkour, Alessandro Di Cerbo, and Claudia Zizzadoro. "Effect of parsley (*Petroselinum crispum*) oil as feed additive on broiler performance, carcass, liver and kidney functions, antioxidant, lipid profile, and immunity." *Animal Science Journal* 95, no. 1 (2024): e13981.
33. Abdelnour, Sameh A., Ibrahim T. El-Ratel, Soliman I. Peris, Ali Ali El-Raghi, and Sara F. Fouda. "Effects of dietary thyme essential oil on blood haematobiochemical, redox status, immunological and reproductive variables of rabbit does exposed to high environmental temperature." *Italian Journal of Animal Science* 21, no. 1 (2022): 51-61.
34. Abdelnour, Sameh A., Mohamed GE Metwally, Laila B. Bahgat, and Mohammed AE Naiel. "Pumpkin seed oil-supplemented diets promoted the growth productivity, antioxidative capacity, and immune response in heat-stressed growing rabbits." *Tropical Animal Health and Production* 55, no. 1 (2023): 55.
35. Singh, A. S., J. O. Alagbe, S. Sharma, R. A. Oluwafemi, and O. C. P. Agubosi. "Effect of dietary supplementation of melon (*Citrullus lanatus*) seed oil on the growth performance and antioxidant status of growing rabbits." *Indonesian Journal of Innovation and Applied Sciences (IJIAS)* 1, no. 2 (2021): 134-143.
36. Alagbe, Olujimi John, and D. N. Anorue. "BIOACTIVE PROFILING OF ESSENTIAL OIL OF *TERMINALIA ARJUNA* STEM BARK COLLECTED FROM ORATHUR VILLAGE, TAMILNADU, INDIA." (2024).

37. Omokore, E. O., and J. O. Alagbe. "Efficacy of dried *Phyllanthus amarus* leaf meal as an herbal feed additive on the growth performance, haematology and serum biochemistry of growing rabbits." *International Journal of Academic Research and Development* 4, no. 3 (2019): 97-104.
38. El-Essawy, Abeer M., U. Y. Anele, A. M. Abdel-Wahed, Ahlam R. Abdou, and I. M. Khattab. "Effects of anise, clove and thyme essential oils supplementation on rumen fermentation, blood metabolites, milk yield and milk composition in lactating goats." *Animal Feed Science and Technology* 271 (2021): 114760.
39. Abou-Elkhair, R., H. A. Ahmed, and S. Selim. "Effects of black pepper (*Piper nigrum*), turmeric powder (*Curcuma longa*) and coriander seeds (*Coriandrum sativum*) and their combinations as feed additives on growth performance, carcass traits, some blood parameters and humoral immune response of broiler chickens." *Asian-Australasian Journal of Animal Sciences* 27, no. 6 (2014): 847.
40. Valiollahi, Mohammad Reza, Yaser Rahimian, Ali Rafiee, and Yasamin Miri. "Effect use ginger (*Zingiber officinale*), black pepper (*Piper nigrum* L) powders on performance, some blood parameters and antibody titer against new castle vaccine on broiler chicks." (2014): 61-66.
41. Costa, L. B., V. V. Almeida, B. Berenchein, M. L. P. Tse, A. C. Andrade, and V. S. Miyada. "Phytobiotic additives and sodium butyrate as alternatives to antibiotics for weanling pigs." *Archivos de zootecnia* 60, no. 231 (2011): 733-744.
42. Volek, Z., T. A. Ebeid, and L. Uhlířová. "The impact of substituting soybean meal and sunflower meal with a mixture of white lupine seeds and rapeseed meal on rabbit doe milk yield and composition, and the growth performance and carcass traits of their litters." *Animal feed science and technology* 236 (2018): 187-195.
43. Shittu, M. T., John Olujimi Alagbe, Olusegun Ojeniyi Ojebiyi, Taiwo Kayode Ojediran, and Trimisiyu Adewale Rafiu. "Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of *Polyalthia longifolia* leaf extract in place of conventional antibiotics." *Animal Science and Genetics* 18, no. 2 (2022).
44. Nwaopara, A. O., M. A. C. Odiye, U. Inegbenebor, and M. I. Aduye. "The combined effects of excessive consumption of ginger, clove, red pepper and black pepper on the histology of the liver." *Pakistan Journal of nutrition* 6, no. 6 (2007): 524-527.
45. Madkour, Mahmoud, Mohamad M. Aboelenin, Eman Younis, Mohamed A. Mohamed, Hussein Hassan, Mahmoud Alagawany, and Mohamed Shourrap. "Hepatic acute-phase response, antioxidant biomarkers and DNA fragmentation of two rabbit breeds subjected to acute heat stress." *Italian Journal of Animal Science* 19, no. 1 (2020): 1568-1576.
46. Sheiha, Asmaa M., Sameh A. Abdelnour, Mohamed E. Abd El-Hack, Asmaa F. Khafaga, Khaled A. Metwally, Jamaan S. Ajarem, Saleh N. Maooda, Ahmed A. Allam, and Mohamed T. El-Saadony. "Effects of dietary biological or chemical-synthesized nano-selenium supplementation on growing rabbits exposed to thermal stress." *Animals* 10, no. 3 (2020): 430.
47. Selim, Shaimaa, Mahmoud F. Seleiman, Mohamed M. Hassan, Ahmed A. Saleh, and Mohamed A. Mousa. "Impact of dietary supplementation with *Moringa oleifera* leaves on performance, meat characteristics, oxidative stability, and fatty acid profile in growing rabbits." *Animals* 11, no. 2 (2021): 248.
48. Abdelnour, Sameh A., Naif A. Al-Gabri, Nesrein M. Hashem, and Antonio Gonzalez-Bulnes. "Supplementation with proline improves haemato-biochemical and reproductive indicators in male rabbits affected by environmental heat-stress." *Animals* 11, no. 2 (2021): 373.
49. Alagbe, J.O., Shittu, M.D., Ojediran, T.K., Omokore, E.A., Oluwayinka, E.O., Anorue, D.N., Bamigboye, S and Effiong, E. (2024). Dietary supplementation of *Ocotea aciphylla* essential oil: effects on growth performance and nutrient digestibility of weaned pigs. *World Journal of Agriculture and Forestry Sciences*, 2(2): 43-49.
50. Alagbe, J. O. "Analysis of bioactive compounds in ethanolic extract of *Xylopiya aethiopyca* leaves using gas chromatography and mass spectrometry (GC-MS) technique." *Journal of Pharmacy and Drug Development* 2, no. 2 (2023): 2836-2322.

Citation: Alagbe J.O., et.al. Toxicological Assessment of Supplemental *Zanthoxylum acanthopodium* essential oil in Rabbits: Evidence from Haemato-biochemical and Histopathological Indices. *Biomed. Biotechnol.Sci.* Vol. 2 Iss. 1. (2026) DOI: 10.58489/2833-0951.009